

Professional Dosing Information for Infants and Children



			Fever Reducer/Pain Reliever						Cold and Flu Remedies							Allergy		Gas Relief																			
SYMPTOMS			Fever Pain	Fever Pain	Fever Pain	Fever Pain	Fever Pain	Fever, Sore Throat Cough, Stuffy Nose Sneezing Runny Nose	Fever, Sore Throat Cough Sneezing Runny Nose	Fever Sore Throat Cough	Fever, Sore Throat Cough, Stuffy Nose Sneezing Runny Nose	Cough Chest Congestion Thins & Loosens Mucus	Stuffy Nose	Cough Nasal Congestion	Stuffy Nose, Itchy Throat, Cough	Itchy, Watery Eyes Itchy Throat/Nose Cough, Sneezing Runny Nose	Itchy, Watery Eyes Itchy Throat/Nose Sneezing Runny Nose	Gas																			
ACTIVE INGREDIENTS			Per 5 mL (1 tsp): Acetaminophen 160 mg	Per 5 mL (1 tsp): Acetaminophen 160 mg	Per 5 mL (1 tsp): Acetaminophen 160 mg	Per 5 mL (1 tsp): Acetaminophen 160 mg	Per 1.25 mL: Ibuprofen 50 mg	Per 5.0 mL (1tsp): Ibuprofen 100 mg	Per 5 mL (1 tsp): Acetaminophen 160 mg, Chlorpheniramine maleate 1 mg, Dextromethorphan HBr 5 mg, Phenylephrine HCl 2.5 mg	Per 5 mL (1 tsp): Acetaminophen 160 mg, Chlorpheniramine maleate 1 mg, Dextromethorphan HBr 5 mg	Per 5 mL (1 tsp): Acetaminophen 160 mg, Chlorpheniramine maleate 1 mg, Dextromethorphan HBr 5 mg, Phenylephrine HCl 2.5 mg	Per 5 mL (1 tsp): Dextromethorphan HBr 5 mg, Guaifenesin, USP 100 mg	Per 5 mL (1 tsp): Phenylephrine HCl 2.5 mg	Per 5 mL (1 tsp): Dextromethorphan HBr 5 mg, Phenylephrine HCl 2.5 mg	Per 5 mL (1 tsp): Diphenhydramine HCl 6.25 mg, Phenylephrine HCl 2.5 mg	Per 5 mL (1 tsp): Diphenhydramine HCl 12.5 mg	Per 5 mL (1 tsp): Cetirizine Hydrochloride 5 mg	Per 0.3 mL: Simethicone 20 mg																			
DOSING GUIDELINES	Weight	Age																																			
	6-11 lbs	0-5 mos	Ask a Doctor	Use PediaCare® Acetaminophen Infants	Use PediaCare® Acetaminophen Infants	Use PediaCare® Acetaminophen Infants	Ask a Doctor	Use PediaCare® Infant IB Ibuprofen Concentrated Oral Suspension	Not Recommended								Ask a Doctor	0.3 mL																			
	12-17 lbs	6-11 mos	Ask a Doctor						1.25 mL	Ask a Doctor	Ask a Doctor	Ask a Doctor	Ask a Doctor	2.5-5 mL (½-1 tsp)	5 mL (1 tsp)	5 mL (1 tsp)			5 mL (1 tsp)	Ask a Doctor	0.3 mL																
	18-23 lbs	12-23 mos	Ask a Doctor						1.875 mL													Ask a Doctor	Ask a Doctor	Ask a Doctor	Ask a Doctor	2.5-5 mL (½-1 tsp)	5 mL (1 tsp)	5 mL (1 tsp)	5 mL (1 tsp)	Ask a Doctor	0.3 mL						
	24-35 lbs	2-3 yrs	5 mL	5 mL (1 tsp)	Use PediaCare® Acetaminophen Infants or Children's 4oz bottle	Use PediaCare® Acetaminophen Infants or Children's 4oz bottle	5 mL	5 mL (1 tsp)	10 mL (2 tsp)	10 mL (2 tsp)	10 mL (2 tsp)	10 mL (2 tsp)	10 mL (2 tsp)	10 mL (2 tsp)	5-10 mL (1-2 tsp)	5 mL (1 tsp) or 10 mL (2 tsp) once daily depending upon severity of symptoms: do not take more than 10 mL (2 tsp) in 24 hours.	0.6 mL																				
	36-47 lbs	4-5 yrs	Use PediaCare® Children Acetaminophen Oral Suspension	7.5 mL (1½ tsp)	One Packet	Use 7.5 mL Packet	7.5 mL											10 mL (2 tsp)	10 mL (2 tsp)	10 mL (2 tsp)	10 mL (2 tsp)											10 mL (2 tsp)	10 mL (2 tsp)	10 mL (2 tsp)	10 mL (2 tsp)	5-10 mL (1-2 tsp)	0.6 mL
	48-59 lbs	6-8 yrs		10 mL (2 tsp)	Use 10 mL Packet	One Packet	10 mL											10 mL (2 tsp)	10 mL (2 tsp)	10 mL (2 tsp)	10 mL (2 tsp)											10 mL (2 tsp)	10 mL (2 tsp)	10 mL (2 tsp)	10 mL (2 tsp)	5-10 mL (1-2 tsp)	0.6 mL
	60-71 lbs	9-10 yrs		12.5 mL (2½ tsp)	Use PediaCare® Children's 4oz bottle	Use PediaCare® Children's 4oz bottle	12.5 mL											10 mL (2 tsp)	10 mL (2 tsp)	10 mL (2 tsp)	10 mL (2 tsp)											10 mL (2 tsp)	10 mL (2 tsp)	10 mL (2 tsp)	10 mL (2 tsp)	5-10 mL (1-2 tsp)	0.6 mL
72-95 lbs	11 yrs	15 mL (3 tsp)		Use PediaCare® Children's 4oz bottle	Use PediaCare® Children's 4oz bottle	15 mL	10 mL (2 tsp)											10 mL (2 tsp)	10 mL (2 tsp)	10 mL (2 tsp)	10 mL (2 tsp)	10 mL (2 tsp)	10 mL (2 tsp)	10 mL (2 tsp)	5-10 mL (1-2 tsp)	0.6 mL											
DOSING FREQUENCY			every 4 hours/ no more than 5 doses daily	every 4 hours/ no more than 5 doses daily	every 4 hours/ no more than 5 doses daily	every 4 hours/ no more than 5 doses daily	every 6-8 hours/ no more than 4 doses daily	every 6-8 hours/ no more than 4 doses daily	every 4 hours/ no more than 5 doses daily	every 4 hours/ no more than 5 doses daily	every 4 hours/ no more than 5 doses daily	every 4 hours/ no more than 5 doses daily	every 4 hours/ no more than 6 doses daily	every 4 hours/ no more than 6 doses daily	every 4 hours/ no more than 6 doses daily	every 4 hours/ no more than 6 doses daily	every 4 hours/ no more than 6 doses daily	refer to dosing guidelines	do not exceed more than 12 doses per day																		
FLAVORS			Cherry Non-Staining Cherry Bubble Gum Grape	Cherry Non-Staining Cherry Grape	Cherry	Cherry	Dye-Free Berry	Dye-Free Berry	Bubble Gum	Cherry	Cherry	Grape	Cherry	Raspberry	Grape	Grape	Cherry Bubble Gum	Dye-Free Grape	Original and Dye-Free																		

Important Instructions for Proper Use: Read and follow the label instructions on all PediaCare® products • Keep all medicines out of the reach of children • Do NOT administer adult medicines to children • If you have any questions, call 1-888-474-3099 weekdays from 8:00 am - 8:00 pm (EST)
 Note: Use weight to dose when possible, otherwise use age. Ideally, the child should be weighed before giving PediaCare® products. **Healthcare professionals should be consulted for dosing in children under the age of two and in children under the age of six for certain cough, cold and allergy medicines.**